



RATINGS SCALE

1.0 Excellent	2.0 Good	3.0 Average	4.0 Fair	5.0 Not Recommended
" " Insufficient data				

Soybean Cyst Nematode (SCN) Resistant: Varieties containing these genes are resistant to the following races of Soybean Cyst Nematode:

F= PI88788 3,6,8,9,10,12,13,14

P= PI548402 1,3,5,6,7,8,10,15

Phytophthora Root Rot Race Resistance: Resistant varieties carry the major gene reported to be resistant to these races:

Rps1-a: 1, 2, 10, 11, 12, 15-18, 24, 26, 27

Rps1-c: 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26

Rps1-k: 1-11, 13-15, 17, 18, 21, 22, 24, 26

Rps3-a: 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25

Rps6: 1-4, 10, 12, 14-16, 18-21, 25

Brown Stem Rot: NG = No Gene

Plant Height: M = Medium, MT = Medium Tall, T = Tall

Plant Type: M= Medium, MB = Medium Bush, B= Bush

Colors: BF= Buff, BL= Black, BR= Brown, G= Gray,

IB= Imperfect Black, P= Purple, W= White

POSITIONING & MANAGEMENT

Dual Phytophthora Root Rot genes and excellent emergence make L 1236 E3 the top choice for early plantings. The medium-tall plant has superior stress tolerance for a wide range of soil types and environments. The dense canopy leads to average Sclerotinia White Mold tolerance. Excellent Iron Deficiency Chlorosis makes it a key line for South Dakota and Minnesota



Highly Productive & Irrigated Fields	X
Moderately Productive/Average Fields	X
Less Productive/Stressed Fields	X

SOYBEAN ADVANTAGES

- Dual PRR genes with superior IDC tolerance
- Excellent emergence and stress tolerance
- Medium-tall plant with excellent standability
- Use caution in areas prone to SWM

PLANT CHARACTERISTICS

Standability	2.0	Pubescence Color	G
Plant Height	MT	Pod Color	T
Plant Type	M	Hilum Color	IB
Flower Color	P		

DISEASE CHARACTERISTICS

Phytophthora Root Rot	1c/3a, 1.5	Sudden Death	3.0
Brown Stem Rot		Frogeye	
White Mold	3.0	Charcoal Rot	

DEFENSIVE RATINGS

SCN Resistance	PI 88788
Iron Chlorosis	2.0
Stress Tolerance	1.5

PLACEMENT

Preferred Row Spacing	
Soil Type	
No-Till Rating	

